Immunoconjugates and humanized antibodies specific for B-cell lymphoma and leukemia cells

Publication number: JP10505231 (T) **Publication date:** 1998-05-26

Inventor(s):
Applicant(s):
Classification:

- international: C12N15/09; A61K39/395; A61K47/48; A61K49/00;

A61P35/02; C07K16/30; C12P21/02; C12P21/08; A61K38/00; C12R1/91; C12N15/09; A61K39/395; A61K47/48; A61K49/00; A61P35/00; C07K16/18; C12P21/02; C12P21/08; A61K38/00; (IPC1-7): C12N15/09; A61K39/395; A61K49/00; C07K16/30;

C12P21/02; C12P21/08; C12P21/02; C12R1/91

- **European:** A61K47/48T4B30; A61K47/48T4B30M; A61K47/48T4B46B;

C07K16/30M

Application number: JP19950507371T 19950811

Priority number(s): WO1995US09641 19950811; US19940289576 19940812

Abstract not available for JP 10505231 (T)

Abstract of corresponding document: US 5789554 (A)

A chimeric LL2 monoclonal antibody is described in which the complementarity determining regions (CDRs) of the light and heavy chains of the murine LL2 anti-B-lymphoma, anti-leukemia cell monoclonal antibody has been recombinantly joined to the human kappa and IgG1 constant region domains, respectively, which retains the immunospecificity and B-cell lymphoma and leukemia cell internalization capacity of the parental murine LL2 monoclonal antibody, and which has the potential of exhibiting reduced human anti-mouse antibody production activity. A humanized LL2 monoclonal antibody is described in which the CDRs of the light and heavy chains have been recombinantly joined to a framework sequence of human light and heavy chains variable regions, respectively, and subsequently linked to human kappa and IgG1 constant region domains, respectively, which retains the immunospecificity and B-lymphoma and leukemia cell internalization capacities of the parental murine and chimeric LL2 monoclonal antibodies, and which has the potential for exhibiting reduced human anti-mouse antibody production activity. Vectors for producing recombinant chimeric and humanized chimeric monoclonal antibodies are provided. Isolated DNAs encoding the amino acid sequences of the LL2 variable light and heavy chain and CDR framework regions are described. Conjugates of chimeric and humanized chimeric LL2 antibodies with cytotoxic agents or labels find use in therapy and diagnosis of B-cell lymphomas and leukemias.

Data supplied from the **esp@cenet** database — Worldwide

Also published as:

US5789554 (A)

WO9604925 (A1)

L114909 (A)

ES2251723 (T3)

more >>